


# Classical Hall effect: resistivity tensor



$$\mathbf{E} = \rho \cdot \mathbf{J}$$

We have shown that:

$$\begin{pmatrix} E_x \\ E_y \end{pmatrix} = \frac{1}{\sigma_0} \begin{pmatrix} 1 & +\omega_c^e \tau_e \\ -\omega_c^e \tau_e & 1 \end{pmatrix} \begin{pmatrix} J_x \\ J_y \end{pmatrix}$$

$$\rho_{xx} = \rho_0$$

Hall Resistivity:

$$\rho_{yx} = -\rho_0 \omega_c^e \tau_e = \frac{B}{(-e)n}$$

Linear in B!

$$\mu_e = \frac{e\tau_e}{m_e^*}$$

$$\omega_c^e = \frac{e \cdot B}{m_e^*}$$

$$\text{For } J_y = 0$$

we get

$$\rho_{xx} = \frac{E_x}{J_x} \quad \rho_{yx} = \frac{E_y}{J_x}$$


# The (integer) quantum Hall effect.




Klaus von Klitzing



1985



K. v. Klitzing, G. Dorda, M. Pepper,  
*Phys. Rev. Lett.* 45, 494 (1980)



Plateaus in  $\rho_{xy}$  vs  $B$ !

[https://www3.physnet.uni-hamburg.de/institute/IAP/Group\\_N/e/semiconductors/semiconductors\\_print.html](https://www3.physnet.uni-hamburg.de/institute/IAP/Group_N/e/semiconductors/semiconductors_print.html)

# The (integer) quantum Hall effect.



Klaus von Klitzing



1985

PRL Milestone

Access |

## New Method for High-Accuracy Determination of the Fine-Structure Constant Based on Quantized Hall Resistance

K. v. Klitzing, G. Dorda, and M. Pepper

Phys. Rev. Lett. **45**, 494 – Published 11 August 1980

An article within the collection: [Letters from the Past - A PRL Retrospective](#)

Citing Articles (2,782)


PDF

Export Citation



Foz do Iguaçú 2015

# Quantum Hall effect: resistivity plateaus



Zeeman splitting of Landau levels: one spin per state

$$e = 1,6 \times 10^{-19} \text{ Coulomb}$$

$$\hbar = 6,58212 \times 10^{-16} \text{ eV.s}$$

“Doubly filled”  $j$ th Landau level (given density  $n$ ):

$$n = j_{\max} \frac{2eB}{h}$$

“Filling factor”  $\nu$  (per spin):

$$\nu = \frac{nh}{eB}$$

**Hall Resistivity:**

$$\rho_{yx} = \frac{B}{(-e)n} = \frac{1}{\nu} \frac{h}{e^2}$$

Resistivity quantization!

$$\frac{h}{e^2} = ??? \quad \Omega$$

Tarefa 16 (em sala, agora)

# The (integer) quantum Hall effect.


Séminaire Poincaré 2 (2004) 1 – 16





Klaus von Klitzing



1985



K. v. Klitzing, G. Dorda, M. Pepper,  
Phys. Rev. Lett. 45, 494 (1980)



Notes 4./5.2.1980

$$E_H = R_s \cdot D \cdot i = \frac{1}{n \cdot e} \cdot B \cdot \frac{I}{b}$$

$$U_H = \frac{B}{n \cdot e} \cdot \frac{I}{b}$$

$$U_H = \frac{e \cdot h \cdot B \cdot I}{e \cdot e \cdot b}$$

$$= \frac{h}{e^2} \cdot I$$

25.76 hV  
25.813


$$N = \frac{eB}{2\pi k} \quad (g_S \cdot g_V = 1)$$

$$\frac{h}{e^2} =$$

$$S_{AB} = \frac{\alpha}{2} \cdot \sqrt{\frac{c}{c_0}}$$

$$\Rightarrow 25.813 \text{ hV}$$

# Quantum Hall effect: resistivity plateaus




Plateaus in the Hall resistivity for integer  $v$

$$\rho_{yx} = -\frac{1}{\nu} \frac{h}{e^2}$$

$$\nu = \frac{nh}{eB}$$

Now, the longitudinal resistivity goes to zero!

$$\rho_{xx} = \frac{\sigma_{xx}}{\sigma_{xx}^2 + \sigma_{xy}^2} \approx \frac{\sigma_{xx}}{\sigma_{xy}^2} \rightarrow 0$$





No magnetoresistance for  $\omega_c \tau$  large ?!?

$$n = j_{\max} \frac{2eB}{h}$$

# Landau edge states

<http://www.sp.phy.cam.ac.uk/research/fundamentals-of-low-dimensional-semiconductor-systems/lowD>



Potential  $V(y)$ : edge states




Fig 4: Red is the graph of hall resistance in units of  $h/e^2$  versus magnetic field in tesla.

