


# Tarefa 19: Haldane model



$$\frac{\mathcal{H}_{\mathbf{q}}}{N} = \begin{pmatrix} M + 2t_2 f(\mathbf{q}, \phi) & t_1 \gamma_{\mathbf{q}} \\ t_1 \gamma_{\mathbf{q}}^* & -M + 2t_2 f(\mathbf{q}, -\phi) \end{pmatrix}$$

$$\gamma_{\mathbf{q}} = 1 + e^{i\mathbf{q} \cdot \mathbf{a}_2} + e^{i\mathbf{q} \cdot (\mathbf{a}_2 - \mathbf{a}_1)}$$

$$f(\mathbf{q}, \phi) = \cos(\mathbf{q} \cdot \mathbf{a}_1 + \phi) + \cos(\mathbf{q} \cdot \mathbf{a}_2 - \phi) + \cos(\mathbf{q} \cdot (\mathbf{a}_2 - \mathbf{a}_1) + \phi)$$

Consider:  $t_1=1$ ,  $\phi=\pi/2$ , and  $\mathbf{a}_1$  and  $\mathbf{a}_2$  as in the left.

- 1) Calculate the Hamiltonian matrix for the Brillouin zone vertices  $\mathbf{q}=\mathbf{K}$  and  $\mathbf{q}=\mathbf{K}'$ . (remember Lista 03!)
- 2) Show that the gap *vanishes* for

$$t_2 = \pm M / (3\sqrt{3})$$

but not in  $\mathbf{K}$  and  $\mathbf{K}'$  at the same time!